

Initial Date: 3/23/2018

Revised Date: 05/23/2023

Section: 2-13

## ***Bleeding Control***

### **Indications:**

Patients with significant traumatic or non-traumatic (i.e., hemodialysis access) external hemorrhage

1. Follow **General Pre-hospital Care-Treatment Protocol** and **Soft Tissue & Orthopedic Injuries-Treatment Protocol**.
2. Apply direct pressure to the wound with clean gauze using universal precautions.
3. If the bleeding is not controlled with direct pressure, treat according to the location of the wound.
  - a. Extremity bleeding - apply tourniquet:(Refer to **Tourniquet Application-Procedure Protocol**)
    - i. If tourniquet unsuccessful apply second/adjacent tourniquet per **Tourniquet Application-Procedure Protocol**.
    - ii. NOTE- tourniquet may be painful, see **Pain Management-Procedure Protocol**.
  - b. Neck, axilla/shoulder or groin bleeding:
    - i. Pack wound with MCA approved hemostatic dressing (if available, following manufacturer's instructions) or clean gauze.
    - ii. Use as much of the dressing/gauze as needed to stop the blood flow.
    - iii. Quickly apply pressure until the bleeding stops. (Approximately 3-5 minutes)
    - iv. Leave the dressing in place and wrap area with bandaging to secure the dressing.
4. Do not remove the bandage or hemostatic dressing/gauze
5. Elevate the injury, if possible.
6. Reassess for bleeding through or around the dressing.
7. For patients who have signs or symptoms of shock, secondary to hemorrhage, refer to **Hemorrhagic Shock-Treatment Protocol**.
8. Transport according to **Adult and Pediatric Trauma Triage-Treatment Protocol** and MCA Transport Protocol

### **Notes:**



If hemostatic dressing is used, contact medical control to advise of application, document time of use, and send packaging from dressing to hospital with patient for removal instructions.